

ULIRGS: ORIGIN AND EVOLUTION¹

I. Márquez and J. Masegosa

Instituto de Astrofísica de Andalucía (CSIC), Granada (Spain)

RESUMEN

Presentamos los resultados preliminares sobre un estudio de la banda de alta luminosidad ($L_{FIR} > 10^{12.3} L_{\odot}$) de las ULIRGs de la muestra de 1Jy. Imágenes profundas, de alta resolución y campo de 7×7 minutos para 2/3 de la muestra indican que las ULIRGs se encuentran generalmente en entornos muy ricos, que se parecen a lo que se espera en los últimos estadios de la fusión de galaxias (encontramos colas de marea largas en sólo 3 sistemas) y todas ellas tienen núcleos múltiples (son dobles en 11 ULIRGs: 2 HII, 4 LINERs, 4 Sy2 y 1 Sy1). Los instrumentos de primera luz previstos para GTC son especialmente indicados para el estudio de la fuente de energía, y permitirán también obtener una visión más clara de la secuencia evolutiva propuesta.

ABSTRACT

We present our preliminary results on the study of the high luminosity ULIRGs ($L_{FIR} > 10^{12.3} L_{\odot}$) in the 1Jy sample. Deep, high resolution optical R images with $7 \text{ arcmin} \times 7 \text{ arcmin}$ field of view for 2/3 of the whole sample, show that they are generally found in very rich environments, they seem to be late stage merger products (long tidal tails are found in only 3 systems) and all of them show multiple nuclei (double nuclei in 11: 2HII, 4 LINERs, 4 Sy2 and 1 Sy1). GTC first light instruments are very well suited for addressing the question of the energy source of ULIRGs, together with a more refined view of the proposed evolutionary sequence.

Key Words: **GALAXIES: ACTIVE, INTERACTIONS, INFRARED**

1. INTRODUCTION

The most conspicuous finding by IRAS (Soifer et al. 1987) was the discovery of a new population of galaxies emitting at FIR wavelengths. Their energies are comparable to those of the most luminous quasars. Such similar power between QSO and luminous infrared galaxies could be explained considering the latter as dust-enshrouded QSOs. The immediate question to answer is the nature of the energy source.

The first IRAS data concerned a sample of 9 galaxies, optically very faint ($L_{IR}/L_B > 20$). Only two of those galaxies had been already classified in the optical range, both with very perturbed morphologies: Arp 220 and NGC 6240. The most likely explanation for the large observed FIR luminosities ($\approx 10^{12} L_{\odot}$) is that they should host very strong UV emitting sources, very deeply obscured by large amounts of dust; the dust would be heated by the UV radiation and re-radiate at FIR. During the late 80's and the beginning of 90's the question whether a black hole or a starburst is the nuclear source was posed. Rowan-Robinson's models for their SED's

showed that, whereas both cirrus and starburst components were always needed, the active nucleus was not invoked in all cases.

Optical spectra showed that most of these galaxies resembled better those of LINERs than Seyferts 1 or 2. The characterization of a complete sample of ULIRGs had to await until 1995: Veilleux et al. (1995); Veilleux et al. (1999) showed that the percentage of HII-like nuclei, LINERs and Seyferts changes as a function of the IR luminosity. For low L_{FIR} most of the nuclei are starburst-like; up to 60% of the most luminous ULIRGs are Seyfert-like, only 20% of them being HII-like. Interestingly, the percentage of LINERs (30%) remains constant in spite of their L_{FIR} level. An evolutionary scheme appears to explain this dependence on L_{FIR} of the nature of the energy source.

2. AN EVOLUTIONARY SEQUENCE

Strongly interacting systems are highly frequent among ULIRGs, and their properties are therefore studied in connection with the effects of merging processes. Sanders et al. (1988) suggested that ULIRGs would evolve in luminosity as precursors of optically selected quasars. Following this scenario, massive star formation induced by strong gravitational inter-

¹The data presented here have been taken using ALFOSC, which is owned by the Instituto de Astrofísica de Andalucía (IAA) and operated at the Nordic Optical Telescope under agreement between IAA and the NBIfAFG of the Astronomical Observatory of Copenhagen

action would be the first energy source; later in the merger sequence more and more star formation is produced and eventually very compact stellar clusters are formed, that could give rise to the formation of a massive black hole (BH); the BH should be fed up with material from the circumnuclear regions, where starbursting processes would take place and consequently an obscured QSO could be observed.

The morphologies of ULIRGS have been classified (Surace et al. 1998; 2000; Surace & Sanders 1999) as: I. Pre-contact; II. First contact without tidal tails; Pre-merger with tidal tails and double nuclei separated (a) more than 10 kpc and (b) less than 10 kpc; IV. Merger with long tidal tails and a single (a) diffuse and (b) compact nucleus; V. Final stage merger, no tidal tails and strong central perturbations.

Following this classification, Veilleux et al. (2001) has found that the percentage of HII-like, LINER and Seyfert (S1, S2) nuclei is: III: 40% HII, 40% LINER, 20% Sy2; IV: 25% HII, 38% LINER, 43% Sy2, 15% Sy1; V: 25% HII, 35% LINER, 40% Sy2, 15% Sy1. That is to say, more advanced mergers host more AGNs (LINERs are constant).

3. OUR AIMS AND PRELIMINARY RESULTS

We plan to characterize the environmental status together with the merger stage, paying attention not only to the very central regions but also to large scale structures that would allow a better determination of the “age” in the expected merger sequence. To do so, high resolution and large field of view optical imaging is being obtained. First, we are in the process of getting R-band images of the whole high luminosity strip ($L_{FIR} > 10^{12.3} L_{\odot}$, 30 galaxies). Among the 19 galaxies observed up to now (2 HII-like, 8 LINERs, 4 Sy2, 5 Sy 1, see the figure for one example) we obtain the following, preliminary conclusions: (1) They are generally found in very rich environments; (2) they seem to be late stage merger products (long tidal tails are found in only 3 systems); (3) all of them show multiple nuclei (double nuclei in 11: 2HII, 4 LINERs, 4 Sy2 and 1 Sy1).

4. ULIRGS WITH GTC

Both the resolution and collecting power of a 10m-class telescope will allow to study the properties of ULIRGs from our nearby Universe to higher redshifts. The question of the nature of the energy source will be obviously better addressed, together with a more refined view of the proposed evolutionary sequence from one type of object to the other, and the study of the nature of the precursors of to-




Fig. 1. R band image of IRAS 07598+6508 (FWHM = 0.8 arcsec). The contour corresponds to $\mu_R = 25.36$ mag/arcsec². Ticks are in arcseconds from the source.

day’s ULIRGs. First light instruments are very well suited for these purposes, both OSIRIS, to get optical to NIR imaging or spectroscopy, and CANARICAM, to get the valuable information provided at MIR wavelengths (note that ULIRGs are one of the scientific cases for CANARICAM). The determination of the nature of the energy source will be completed with the advent of EMIR.

We acknowledge financial support from the Spanish Ministerio de Ciencia y Tecnología through the programs PB98-0521 and AYA2001-2089.

REFERENCES

- Sanders, D.B., Soifer, B.T., Scoville, N.Z. & Sargent, A.I. 1988, ApJ 324, 55
- Soifer, B.T., Sanders, D.B., Madore, B.F., Neugebauer, G., Danielson, G.E. et al. 1987, ApJ 320, 238
- Surace, J.A., Sanders, D.B., Vacca, W.D., Veilleux, S. & Mazzarella, J.M. 1998, ApJ 492, 116
- Surace, J.A. & Sanders, D.B. 1999, ApJ 512, 162
- Surace, J.A., Sanders, D.B. & Evans, A.S. 2000, ApJ 529, 170
- Veilleux, S., Kim, D.-C., Sanders, D.B., Mazzarella & J.M., Soifer, B.T. 1995, ApJS 98, 171
- Veilleux, S., Kim, D.-C. & Sanders, D.B. 1999, ApJ 522, 113
- Veilleux, S., Kim, D.-C. & Sanders, D.B. 2001, in “QSO Hosts and Their Environments”, Ed. by I. Márquez et al., Kluwer Academic/Plenum Publishers, p. 165

I. Márquez and J. Masegosa: Instituto de Astrofísica de Andalucía (CSIC), Apdo 3004, 18080 Granada (Spain)
(isabel@iaa.es, pepa@iaa.es).